arXiv:1210.2539v1 [math.DG] 9 Oct 2012

The mean curvature flow for invariant hypersurfaces in a Hilbert space with a free group action

Naoyuki Koike

Abstract

In this paper, we study the regularized mean curvature flow starting from a G-invariant hypersurfaces in a Hilbert space equipped with an isometric and free action of Hilbert Lie group G whose orbits are minimal regularizable submanifolds. First we obtain the evolution equations for some geometric quantities along the regularized mean curvature flow. Next, by using the evolution equations, we prove a strictly horizontally convexity preservability theorem for the regularized mean curvature flow.

1 Introduction

R. S. Hamilton ([Ha]) proved the existenceness and the uniqueness (in short time) of solutions satisfying any initial condition of a weakly parabolic equation for sections of a finite dimensional vector bundle. The Ricci flow equation for Riemannian metrics on a fixed compact manifold M is a weakly parabolic equation, where we note that the Riemannian metrics are sections of the $(0,2)$-tensor bundle $T^{(0,2)}M$ of M. Let f_t $(0 \leq t < T)$ be a C^∞-family of immersions of M into the m-dimensional Euclidean space $\mathbb{R}^m$. Define a map $F : M \times [0,T) \to \mathbb{R}^m$ by $F(x,t) := f_t(x)$ $((x,t) \in M \times [0,T))$. The mean curvature flow equation is described as

$$\frac{\partial F}{\partial t} = \triangle_t f_t,$$

where $\triangle_t$ is the Laplacian operator of the metric g_t on M induced from the Euclidean metric of $\mathbb{R}^m$ by f_t. Here we note that $\triangle_t f_t$ is equal to the mean curvature vector of f_t. This evolution equation is a weakly parabolic equation, where we note that the immersions f_t's are regarded as sections of the trivial bundle $M \times \mathbb{R}^m$ over M under the identification of f_t and its graph immersion $\mathrm{id}_M \times f : M \to M \times \mathbb{R}^m$ (id_M : the identity map of M). Hence we can apply the Hamilton's result to this evolution equation and hence can show the existenceness and the uniqueness (in short time) of solution of this evolution equation satisfying any initial condition. In this paper, we consider the case where the ambient space is a (separable infinite dimensional) Hilbert space V. Let M be a Hilbert manifold and f_t $(0 \leq t < T)$ be a C^∞-family of immersions of M into V. Assume that f_t is regularizable, where "regularizability" means that, for each normal vector v of M, the shape operator A_v is a compact operator, and that the regularized trace $\mathrm{Tr}_r\, A_v$ of A_v and the trace $\mathrm{Tr}\, A_v^2$ of A_v^2 exist. Denote by H_t the regularized mean curvature vector of f_t. See the next section about the definitions of $\mathrm{Tr}_r\, A_v$ and H_t. Define a map $F : M \times [0, T) \to V$ as above in terms of f_t's. We call f_t's $(0 \leq t < T)$ the *regularized mean curvature flow* if the following evolution equation holds:

$$\frac{\partial F}{\partial t} = \triangle_t^r f_t. \tag{1.1}$$

Here $\triangle_t^r f_t$ is defined as the vector field along f_t satisfying

$$\langle \triangle_t^r f_t, v\rangle := \mathrm{Tr}_r \langle (\nabla^t df_t)(\cdot,\cdot), v\rangle^\sharp \quad (\forall\, v \in V),$$

where ∇^t is the Riemannian connection of the metric g_t on M induced from the metric $\langle\ ,\ \rangle$ of V by f_t, $\langle (\nabla^t df_t)(\cdot,\cdot), v\rangle^\sharp$ is the $(1,1)$-tensor field on M defined by $g_t(\langle (\nabla^t df_t)(\cdot,\cdot), v\rangle^\sharp(X), Y) = \langle (\nabla^t df_t)(X,Y), v\rangle$ $(X, Y \in TM)$ and $\mathrm{Tr}_r(\cdot)$ is the regularized trace of $(\cdot)$. Note that $\triangle_t^r f_t$ is equal to H_t. In general, the existenceness and the uniqueness (in short time) of solutions of this evolution equation satisfying any initial condition has not been shown yet. For we cannot apply the Hamilton's result to this evolution equation because it is regarded as the evolution equation for sections of the *infinite* dimensional vector bundle $M \times V$ over M. However we can show the existenceness and the uniqueness (in short time) of solutions of this evolution equation in special case. In this paper, we consider a isometric free action of a Hilbert Lie group G on a Hilbert space V whose orbits are regularized minimal, that is, they are regularizable submanifold and their regularized mean curvature vectors vanish. Let $M(\subset V)$ be a G-invariant submanifold in V whose image by the orbit map of the G-action is compact, and f the inclusion map of M into V. We first show that the regularized mean curvature flow starting from M exists uniquely in short time (see Proposition 3.1). In particular, we consider the case where M is a hypersurface. The first purposes of this paper is to obatin the evolution equations for various geometrical quantities associated with the regularized mean curvature flow starting from such an invariant hypersurface. (see Section 3). The second purpose is to prove a maximum principal for an evolution equation related to a G-invariant symmetric $(0,2)$-tensor fields S_t's on a Hilbert manifold M equipped with an isometric free Hilbert Lie group action G such that M/G is a finite dimensional compact manifold (see Section 4). The third purpose of this paper is to prove a strictly horizontally convexity preservability theorem for the regularized mean curvature flow starting from the above invariant hypersurface by using the evolution equations in Section 3 and imitating the discussion in the proof of a maximal principal in Section 4 (see Section 5). From this theorem, we can derived the strictly convexity preservability theorem for the mean curvature flow in a complete Riemannian manifold by G. Huisken ([Hu2]) in the case where the ambient space is a compact Riemannian reductive homogeneous space whose isometry group is compact and semi-simple.

2 Basic notions and facts

Let f_t $(0 \leq t < T)$ be a one-parameter C^∞-family of immersions of a manifold M into a (finite dimensional) Riemannian manifold N, where T is a positive constant or $T = \infty$. Denote by H_t the mean curvature vector of f_t. Define a map $F : M \times [0,T) \to N$ by $F(x,t) = f_t(x)$ $((x,t) \in M \times [0,T))$. If, for each $t \in [0,T)$, $\dfrac{\partial F}{\partial t} = H_t$ holds, then f_t $(0 \leq t < T)$ is called a *mean curvature flow.*

Let f be an immersion of an (infinite dimensional) Hilbert manifold M into a Hilbert space V and A the shape tensor of f. If $\operatorname{codim} M < \infty$ and A_v is a compact operator for each normal vector v of f, then M is called a *Fredholm submanifold.* In this paper, we then call f a *Fredholm immersion.* Furthermore, if, for each normal vector v of M, the regularized trace $\operatorname{Tr}_r A_v$ and $\operatorname{Tr} A_v^2$ exist, then M is called *regularizable submanifold*, where $\operatorname{Tr}_r A_v$ is defined by $\operatorname{Tr}_r A_v := \sum\limits_{i=1}^{\infty}(\mu_i^+ + \mu_i^-)$ $(\mu_1^- \leq \mu_2^- \leq \cdots \leq 0 \leq \cdots \leq \mu_2^+ \leq \mu_1^+$: the spectrum of $A_v)$. In this paper, we then call f *regularizable immersion.* If f is a regulalizable immersion, then the regularized mean curvature vector H of f is defined by $\langle H, v\rangle = \operatorname{Tr}_r A_v$ $(\forall\, v \in T^\perp M)$, where $\langle\ ,\ \rangle$ is the inner product of V and $T^\perp M$ is the normal bundle of f. In particular, if f is of codimension one, then we call the norm $||H||$ of H the regularized mean curvature function of f.

Let f_t $(0 \leq t < T)$ be a C^∞-family of regularizable immersions of M into V. Denote by H_t the regularized mean curvature vector of f_t. Define a map $F : M \times [0,T) \to V$ by $F(x,t) := f_t(x)$ $((x,t) \in M \times [0,T))$. If $\frac{\partial F}{\partial t} = H_t$ holds, then we call f_t $(0 \leq t < T)$ the *regularized mean curvature flow.* It has not been known whether the regulalized mean curvature flow starting from any regularizable hypersurface exists uniquely in short time. However its existence and uniqueness (in short time) is shown in a special case (see Proposition 3.1).

3 Evolution equations

Let $G \curvearrowright V$ be an isometric free action with minimal regularizable orbit of a Hilbert Lie group G on a Hilbert space V equipped with an inner product $\langle\ ,\ \rangle$. Since each fibre is regularizable, the orbit space V/G is a finite dimensional manifold. Denote by $\phi : V \to V/G$ the orbit map and N the orbit space V/G. Here we give an example of such an isometric free action of a Hilbert Lie group on a Hilbert space.

Example. Let G be a compact semi-simple Lie group and K a closed subgroup of G. Denote by $\mathfrak{g}$ and $\mathfrak{k}$ the Lie algebras of G and K, respectively. Assume that a reductive decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ exists. Let B be the Killing form of $\mathfrak{g}$ and g the bi-invariant metric of G induced from B. Let $H^0([0,1],\mathfrak{g})$ be the Hilbert space of all paths in the Lie algebra $\mathfrak{g}$ of G which are L^2-integrable with respect to B. Also, let $H^1([0,1],G)$ the Hilbert Lie group of all paths in G which are of class H^1 with respect to g. This group $H^1([0,1],G)$ acts on $H^0([0,1],\mathfrak{g})$ isometrically and transitively as a gauge action:

$$\begin{aligned}(g * u)(t) &= \mathrm{Ad}_G(g(t))(u(t)) - (R_{g(t)})_*^{-1}(g'(t)) \\ &\qquad (g \in H^1([0,1],G),\ u \in H^0([0,1],\mathfrak{g})),\end{aligned}$$

where Ad_G is the adjoint representation of G and $R_{g(t)}$ is the right translation by $g(t)$ and g' is the weak derivative of g. Set $P(G, \{e\}\times K) := \{g \in H^1([0,1],G) \,|\, (g(0), g(1)) \in \{e\} \times K\}$, where e is the identity element of G. This group $P(G, \{e\} \times K)$ acts on $H^0([0,1],\mathfrak{g})$ freely and isometrically, and the orbit space of this action is diffeomorphic to G/K. Furthermore, each orbit of this action is a minimal regularizable submanifold.

Give N the Riemannian metric such that ϕ is a Riemannian submersion. Let $M(\subset V)$ be a G-invariant submanifold in V and set $\overline{M} := \phi(M)$. Let f be an inclusion map of M into V and $\overline{f}$ that of $\overline{M}$ into N. Let $\overline{f}_t$ $(0 \le t < T)$ the mean curvature flow starting from $\overline{f}$. According to Lemma 3.2 of [Hu2], $\overline{f}_t$'s $(0 \le t < T)$ are embeddings because $\overline{f}$ is an embedding. Set $\overline{M}_t := \overline{f}_t(\overline{M})$. Define a map $\overline{F} : \overline{M} \times [0,T) \to N$ by $\overline{F}(x,t) := \overline{f}_t(x)$ $((x,t) \in \overline{M} \times [0,T))$. Take $x \in \overline{M}$ and $u \in \phi^{-1}(x)$. Define a curve $c_x : [0,T) \to N$ by $c_x(t) := \overline{f}_t(x)$ and let $(c_x)^L_u : [0,T) \to V$ be the horizontal lift of c_x to u. Define an immersion $f_t : M \hookrightarrow V$ by $f_t(u) = (c_x)^L_u(t)$ $(u \in \widetilde{M})$ and a map $F : M \times [0,T) \to V$ by $F(u,t) = f_t(u)$ $((u,t) \in M \times [0,T))$. Since $\overline{f}_t$'s $(0 \le t < T)$ are embeddings, f_t's $(0 \le t < T)$ also are embeddings. Set $M_t := f_t(M)$.

Proposition 3.1. *The flow f_t $(0 \le t < T)$ is the regularized mean curvature flow starting from f.*

Proof. Denote by $\overline{H}_t$ the mean curvature vector of $\overline{f}_t$ and H_t the regularized mean curvature vector of f_t. It is clear that $\phi\circ f_t=\overline{f}_t\circ\phi_0$. Hence H_t coincides with the horizontal lift $\overline{H}_t^L$ of $\overline{H}_t$. From the definition of F, we have

$$\frac{\partial F}{\partial t}(u,t)=((c_x)_u^L)'(t)=\left(\frac{\partial\overline{F}}{\partial t}(x,t)\right)^L_{(c_x)_u^L(t)}=(\overline{H}_t^L)_u=(H_t)_u.$$

Thus f_t $(0\le t<T)$ is a regularized mean curvature flow. It is clear that $f_0=f$. This completes the proof. q.e.d.

Assume that the codimension of M is equal to one. Denote by $\widetilde{\mathcal{H}}$ (resp. $\widetilde{\mathcal{V}}$) the horizontal (resp. vertical) distribution of ϕ. Denote by $\mathrm{pr}_{\widetilde{\mathcal{H}}}$ (resp. $\mathrm{pr}_{\widetilde{\mathcal{V}}}$) the orthogonal projection of TV onto $\widetilde{\mathcal{H}}$ (resp. $\widetilde{\mathcal{V}}$). For simplicity, for $X\in TV$, we denote $\mathrm{pr}_{\widetilde{\mathcal{H}}}(X)$ (resp. $\mathrm{pr}_{\widetilde{\mathcal{V}}}(X)$) by $X_{\widetilde{\mathcal{H}}}$ (resp. $X_{\widetilde{\mathcal{V}}}$). Define a distribution $\mathcal{H}_t$ on M by $f_{t*}((\mathcal{H}_t)_u)=f_{t*}(T_uM)\cap\widetilde{\mathcal{H}}_{f_t(u)}$ $(u\in M)$ and a distribution $\mathcal{V}_t$ on M by $f_{t*}((\mathcal{V}_t)_u)=f_{t*}(T_uM)\cap\widetilde{\mathcal{V}}_{f_t(u)}$ $(u\in M)$. Note that $\mathcal{V}_t$'s $(t\in[0,T))$ coincide. Let g_t,h_t,A_t,H_t and ξ_t the induced metric, the second fundamental form, the shape tensor and the regularized mean curvature vector and the unit normal vector field of f_t, respectively, and $\overline{g}_t,\overline{h}_t,\overline{A}_t,\overline{H}_t$ and $\overline{\xi}_t$ the corresponding quantities of $\overline{f}_t$. The group G acts on M through f_t. Since $\phi:V\to V/G$ is a G-bundle and $\widetilde{\mathcal{H}}$ is a connection of the bundle, it follows from Proposition 3.1 that this action is independent of the choice of t. It is clear that quantities g_t,h_t,A_t and H_t are G-invariant. Also, let ∇^t (resp. $\overline{\nabla}^t$) the Riemannian connection of g_t (resp. $\overline{g}_t$). Let π_M be the projection of $M\times[0,T)$ onto M and $\pi_{\overline{M}}$ that of $\overline{M}\times[0,T)$ onto $\overline{M}$. For a vector bundle E over M, denote by π_M^*E the induced bundle of E by π_M. Also denote by $\Gamma(E)$ the space of all sections of E. Define a section g of $\pi_M^*(T^{(0,2)}M)$ by $g(u,t)=(g_t)_u$ $((u,t)\in M\times[0,T))$, where $T^{(0,2)}M$ is the $(0,2)$-tensor bundle of M. Similarly, we define a section h of $\pi_M^*(T^{(0,2)}M)$, a section A of $\pi_M^*(T^{(1,1)}M)$, a map $H:M\times[0,T)\to TV$, a map $\xi:M\times[0,T)\to TV$, sections $\overline{g},\overline{h}$ of $\pi_{\overline{M}}^*(T^{(0,2)}\overline{M})$, a section $\overline{A}$ of $\pi_{\overline{M}}^*(T^{(1,1)}\overline{M})$, maps $\overline{H}:\overline{M}\times[0,T)\to TN$ and $\overline{\xi}:\overline{M}\times[0,T)\to TN$. We regard H and ξ V-valued functions over $M\times[0,T)$ under the identification of T_uV's $(u\in V)$ and V. Define a subbundle $\mathcal{H}$ (resp. $\mathcal{V}$) of π_M^*TM by $\mathcal{H}_{(u,t)}:=(\mathcal{H}_t)_u$ (resp. $\mathcal{V}_{(u,t)}:=(\mathcal{V}_t)_u$). Denote by $\mathrm{pr}_{\mathcal{H}}$ (resp. $\mathrm{pr}_{\mathcal{V}}$) the orthogonal projection of $\pi_M^*(TM)$ onto $\mathcal{H}$ (resp. $\mathcal{V}$). For simplicity, for $X\in\pi_M^*(TM)$, we denote $\mathrm{pr}_{\mathcal{H}}(X)$ (resp. $\mathrm{pr}_{\mathcal{V}}(X)$) by $X_{\mathcal{H}}$ (resp. $X_{\mathcal{V}}$). The bundle $\pi_M^*(TM)$ (resp. $\pi_{\overline{M}}^*(T\overline{M})$) is regarded as a subbundle of $T(M\times[0,T))$ (resp. $T(\overline{M}\times[0,T))$). For a section B of $\pi_M^*(T^{(r,s)}M)$, we define $\frac{\partial B}{\partial t}$ by $\left(\frac{\partial B}{\partial t}\right)_{(u,t)}:=\frac{dB_{(u,t)}}{dt}$, where the right-hand side of this relation is the derivative of the vector-valued function $t \mapsto B_{(u,t)} \,(\in T_u^{(r,s)}M)$. Also, we define a section $B_{\mathcal{H}}$ of $\pi_M^*(T^{(r,s)}M)$ by

$$B_{\mathcal{H}} = \underbrace{(\mathrm{pr}_{\mathcal{H}} \otimes \cdots \otimes \mathrm{pr}_{\mathcal{H}})}_{(r-\text{times})} \circ B \circ \underbrace{(\mathrm{pr}_{\mathcal{H}} \otimes \cdots \otimes \mathrm{pr}_{\mathcal{H}})}_{(s-\text{times})}.$$

The restriction of $B_{\mathcal{H}}$ to $\mathcal{H} \times \cdots \times \mathcal{H}$ (s-times) is regararded as a section of the (r,s)-tensor bundle $\mathcal{H}^{(r,s)}$ of $\mathcal{H}$. This restriction also is denoted by the same symbol $B_{\mathcal{H}}$. For a tangent vector field X on M (or an open set U of M), we define a section $\bar{X}$ of π_M^*TM (or $\pi_M^*TM|_U$) by $\bar{X}_{(u,t)} := X_u$ ($(u,t) \in M \times [0,T)$). Denote by $\widetilde{\nabla}$ the Riemannian connection of V. Define a connection ∇ on π_M^*TM by

$$\nabla_X Y := \nabla^t_X Y_{(\cdot,t)} \;\text{ and }\; \nabla_{\frac{\partial}{\partial t}} Y := \frac{dY_{(u,\cdot)}}{dt}$$

for $X \in T_{(u,t)}(M \times \{t\})$ and $Y \in \Gamma(\pi_M^*TM)$, where $\dfrac{dY_{(u,t)}}{dt}$ is the derivative of the vector-valued function $t \mapsto Y_{(u,t)} \,(\in T_uM)$. Define a connection $\nabla^{\mathcal{H}}$ of $\mathcal{H}$ by $\nabla^{\mathcal{H}}_X Y := (\nabla_X Y)_{\mathcal{H}}$ for $X \in T(M \times [0,T))$ and $Y \in \Gamma(\mathcal{H})$. Similarly, define a connection $\nabla^{\mathcal{V}}$ of $\mathcal{V}$ by $\nabla^{\mathcal{V}}_X Y := (\nabla_X Y)_{\mathcal{V}}$ for $X \in T(M \times [0,T))$ and $Y \in \Gamma(\mathcal{V})$. First we shall show the following evolution equation for g.

Lemma 3.2. *The induced metrics $(g_{\mathcal{H}})_t$ satisfies the following evolution equation:*

$$\frac{\partial g_{\mathcal{H}}}{\partial t} = -2||H||h_{\mathcal{H}},$$

where $||H|| := \sqrt{g(H,H)}$.

Proof. Take $X, Y \in \Gamma(TM)$. We have

$$\begin{aligned}
&\frac{\partial g_{\mathcal{H}}}{\partial t}(\bar{X},\bar{Y}) = \frac{\partial}{\partial t} g_{\mathcal{H}}(\bar{X},\bar{Y}) = \frac{\partial}{\partial t} g(\bar{X}_{\mathcal{H}},\bar{Y}_{\mathcal{H}}) = \frac{\partial}{\partial t}\langle F_*\bar{X}_{\mathcal{H}}, F_*\bar{Y}_{\mathcal{H}}\rangle \\
&= \langle \frac{\partial}{\partial t}(\bar{X}_{\mathcal{H}}F), \bar{Y}_{\mathcal{H}}F\rangle + \langle \bar{X}_{\mathcal{H}}F, \frac{\partial}{\partial t}(\bar{Y}_{\mathcal{H}}F)\rangle \\
&= \langle \bar{X}_{\mathcal{H}}\left(\frac{\partial F}{\partial t}\right) + [\frac{\partial}{\partial t}, \bar{X}_{\mathcal{H}}]F, \bar{Y}_{\mathcal{H}}F\rangle + \langle \bar{X}_{\mathcal{H}}F, \bar{Y}_{\mathcal{H}}\left(\frac{\partial F}{\partial t}\right) + [\frac{\partial}{\partial t}, \bar{Y}_{\mathcal{H}}]F\rangle \\
&= \langle \bar{X}_{\mathcal{H}}(||H||\xi), \bar{Y}_{\mathcal{H}}F\rangle + \langle \bar{X}_{\mathcal{H}}F, \bar{Y}_{\mathcal{H}}(||H||\xi)\rangle \\
&= -||H||g(A\bar{X}_{\mathcal{H}},\bar{Y}_{\mathcal{H}}) - ||H||g(\bar{X}_{\mathcal{H}},A\bar{Y}_{\mathcal{H}}) = -2||H||h_{\mathcal{H}}(\bar{X},\bar{Y}),
\end{aligned}$$

where we use $[\frac{\partial}{\partial t}, \bar{X}_{\mathcal{H}}] \in \mathcal{V}$ and $[\frac{\partial}{\partial t}, \bar{Y}_{\mathcal{H}}] \in \mathcal{V}$. Thus we obtain the desired evolution equation. q.e.d.

Next we show the following evolution equation for ξ.

Lemma 3.3. *The unit normal vector fields* ξ_t *satisfies the following evolution equation:*

$$\frac{\partial \xi}{\partial t} = -F_*(\mathrm{grad}_g ||H||),$$

where $\mathrm{grad}_g(||H||)$ *is the element of* $\pi_M^*(TM)$ *defined by* $d||H||(X) = g(\mathrm{grad}_g ||H||, X)$ *for any* $X \in \pi_M^*(TM)$.

Proof. Since $\langle \xi, \xi \rangle = 1$, we have $\langle \frac{\partial \xi}{\partial t}, \xi \rangle = 0$. Hence $\frac{\partial \xi}{\partial t}$ is tangent to $f_t(M)$. Take any $(u_0, t_0) \in M \times [0, T)$. Let $\{e_i\}_{i=1}^{\infty}$ be an orthonormal base of $T_{u_0}M$ with respect to $g_{(u_0,t_0)}$. By the Fourier expanding $\frac{\partial \xi}{\partial t}|_{t=t_0}$, we have

$$\begin{aligned}
&\left.\frac{\partial \xi}{\partial t}\right|_{t=t_0} = \sum \langle \left.\frac{\partial \xi}{\partial t}\right|_{t=t_0}, f_{t_0*}(\bar{e}_i|_{t=t_0})\rangle f_{t_0*}(\bar{e}_i|_{t=t_0}) \\
&= -\sum \langle \xi_{t_0}, \left.\frac{\partial f_{t*}(\bar{e}_i)}{\partial t}\right|_{t=t_0} \rangle f_{t_0*}(\bar{e}_i|_{t=t_0}) \\
&= -\sum \langle \xi_{t_0}, \left.\frac{\partial}{\partial t}(\bar{e}_i F)\right|_{t=t_0} \rangle f_{t_0*}(\bar{e}_i|_{t=t_0}) \\
&= -\sum \langle \xi_{t_0}, \bar{e}_i \left(\left.\frac{\partial F}{\partial t}\right|_{t=t_0} \right) \rangle f_{t_0*}(\bar{e}_i|_{t=t_0}) \\
&= -\sum \langle \xi_{t_0}, (\bar{e}_i H)|_{t=t_0} \rangle f_{t_0*}(\bar{e}_i|_{t=t_0}) \\
&= -\sum (\bar{e}_i ||H||)|_{t=t_0} f_{t_0*}(\bar{e}_i|_{t=t_0}) \\
&= -\sum g_{t_0}(\mathrm{grad}_{g_{t_0}} ||H_{t_0}||, \bar{e}_i|_{t=t_0}) f_{t_0*}(\bar{e}_i|_{t=t_0}) \\
&= -f_{t_0*}(\mathrm{grad}_{g_{t_0}} ||H_{t_0}||) = -F_*(\mathrm{grad}_g ||H||)|_{t=t_0}
\end{aligned}$$

on U, where we use $[\frac{\partial}{\partial t}, \bar{e}_i] = 0$. Here we note that $\sum (\cdot)_i$ means $\lim\limits_{k\to\infty} \sum_{i \in S_k} (\cdot)_i$ as $S_k := \{i \mid |(\cdot)_i| > \frac{1}{k}\}$ $(k \in \mathbb{N})$. In particular, we obtain $(\frac{\partial \xi}{\partial t})_{(u_0,t_0)} = -(F_*((\mathrm{grad}_g ||H||))_{(u_0,t_0)}$. This completes the proof. q.e.d.

Let a C^∞-family S_t $(0 \le t < T)$ of a (r, s)-tensor fields on M and S a section of $\pi_M^*(T^{(r,s)}M)$ defined by $S_{(u,t)} := (S_t)_u$. We define a section $\triangle_{\mathcal{H}} S$ of $\pi_M^*(T^{(r,s)}M)$ by

$$(\triangle_{\mathcal{H}} S)_{(u,t)} := \sum_{i=1}^{n} \nabla_{e_i} \nabla_{e_i} S,$$

where ∇ is the connection of $\pi_M^*(T^{(r,s)}M)$ (or $\pi_M^*(T^{(r,s+1)}M)$) induced from ∇ and $\{e_1, \cdots, e_n\}$ is an orthonormal base of $\mathcal{H}_{(u,t)}$ with respect to $(g_{\mathcal{H}})_{(u,t)}$. Also, we define a section $\bar{\triangle}_{\mathcal{H}}S_{\mathcal{H}}$ of $\mathcal{H}^{(r,s)}$ by

$$(\bar{\triangle}_{\mathcal{H}}S_{\mathcal{H}})_{(u,t)} := \sum_{i=1}^{n} \nabla^{\mathcal{H}}_{e_i}\nabla^{\mathcal{H}}_{e_i}S_{\mathcal{H}},$$

where $\nabla^{\mathcal{H}}$ is the connection of $\mathcal{H}^{(r,s)}$ (or $\mathcal{H}^{(r,s+1)}$) induced from $\nabla^{\mathcal{H}}$ and $\{e_1, \cdots, e_n\}$ is as above. Let $\mathcal{A}^{\phi}$ be the section of $T^*V \otimes T^*V \otimes TV$ defined by

$$\mathcal{A}^{\phi}_X Y := (\widetilde{\nabla}_{X_{\widetilde{\mathcal{H}}}} Y_{\widetilde{\mathcal{H}}})_{\widetilde{\mathcal{V}}} + (\widetilde{\nabla}_{X_{\widetilde{\mathcal{H}}}} Y_{\widetilde{\mathcal{V}}})_{\widetilde{\mathcal{H}}} \quad (X, Y \in TV).$$

Also, let $\mathcal{T}^{\phi}$ be the section of $T^*V \otimes T^*V \otimes TV$ defined by

$$\mathcal{T}^{\phi}_X Y := (\widetilde{\nabla}_{X_{\widetilde{\mathcal{V}}}} Y_{\widetilde{\mathcal{H}}})_{\widetilde{\mathcal{V}}} + (\widetilde{\nabla}_{X_{\widetilde{\mathcal{V}}}} Y_{\widetilde{\mathcal{V}}})_{\widetilde{\mathcal{H}}} \quad (X, Y \in TV).$$

Also, let $\mathcal{A}_t$ be the section of $T^*M \otimes T^*M \otimes TM$ defined by

$$(\mathcal{A}_t)_X Y := (\nabla^t_{X_{\mathcal{H}_t}} Y_{\mathcal{H}_t})_{\mathcal{V}_t} + (\nabla^t_{X_{\mathcal{H}_t}} Y_{\mathcal{V}_t})_{\mathcal{H}_t} \quad (X, Y \in TM).$$

Also let $\mathcal{A}$ be the section of $\pi_M^*(T^*M \otimes T^*M \otimes TM)$ defined in terms of $\mathcal{A}_t$'s ($t \in [0, T)$). Also, let $\mathcal{T}_t$ be the section of $T^*M \otimes T^*M \otimes TM$ defined by

$$(\mathcal{T}_t)_X Y := (\nabla^t_{X_{\mathcal{V}_t}} Y_{\mathcal{V}_t})_{\mathcal{H}_t} + (\nabla^t_{X_{\mathcal{V}_t}} Y_{\mathcal{H}_t})_{\mathcal{V}_t} \quad (X, Y \in TM).$$

Also let $\mathcal{T}$ be the section of $\pi_M^*(T^*M \otimes T^*M \otimes TM)$ defined in terms of $\mathcal{T}_t$'s ($t \in [0, T)$). Clearly we have

$$F_*(\mathcal{A}_X Y) = \mathcal{A}^{\phi}_{F_*X} F_*Y$$

for $X, Y \in \mathcal{H}$ and

$$F_*(\mathcal{T}_W X) = \mathcal{T}^{\phi}_{F_*W} F_*X$$

for $X \in \mathcal{H}$ and $W \in \mathcal{V}$. Let E be a vector bundle over M. For a section S of $\pi_M^*(T^{(0,r)}M \otimes E)$, we define $\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} S(\cdots, \overset{j}{\bullet}, \cdots, \overset{k}{\bullet}, \cdots)$ by

$$(\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} S(\cdots, \overset{j}{\bullet}, \cdots, \overset{k}{\bullet}, \cdots))_{(u,t)} = \sum_{i=1}^{n} S_{(u,t)}(\cdots, \overset{j}{e_i}, \cdots, \overset{k}{e_i}, \cdots)$$

($(u, t) \in M \times [0, T)$), where $\{e_1, \cdots, e_n\}$ is an orthonormal base of $\mathcal{H}_{(u,t)}$ with respect to $(g_{\mathcal{H}})_{(u,t)}$, $S(\cdots, \overset{j}{\bullet}, \cdots, \overset{k}{\bullet}, \cdots)$ means that $\bullet$ is entried into the j-th component and the k-th component of S and $S_{(u,t)}(\cdots, \overset{j}{e_i}, \cdots, \overset{k}{e_i}, \cdots)$ means that e_i is entried into the j-th component and the k-th component of $S_{(u,t)}$.

Then we have the following relation.

Lemma 3.4. *Let S be a section of $\pi_M^*(T^{(0,2)}M)$ which is symmetric with respect to g. Then we have*

$$\begin{aligned}(\triangle_{\mathcal{H}}S)_{\mathcal{H}}(X,Y) = (\bar{\triangle}_{\mathcal{H}}S_{\mathcal{H}})(X,Y)&\\ -2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}((\nabla_{\bullet}S)(\mathcal{A}_{\bullet}X,Y)) - 2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}((\nabla_{\bullet}S)(\mathcal{A}_{\bullet}Y,X))&\\ -\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}S(\mathcal{A}_{\bullet}(\mathcal{A}_{\bullet}X),Y) - \mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}S(\mathcal{A}_{\bullet}(\mathcal{A}_{\bullet}Y),X)&\\ -\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}S((\nabla_{\bullet}\mathcal{A})_{\bullet}X,Y) - \mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}S((\nabla_{\bullet}\mathcal{A})_{\bullet}Y,X)&\\ -2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}S(\mathcal{A}_{\bullet}X,\mathcal{A}_{\bullet}Y)&\end{aligned}$$

for $X,Y \in \mathcal{H}$, where ∇ is the connection of $\pi_M^*(T^{(1,2)}M)$ induced from ∇.

Proof. Take any $(u_0,t_0) \in M \times [0,T)$. Let $\{e_1,\cdots,e_n\}$ be an orthonormal base of $\mathcal{H}_{(u_0,t_0)}$ with respect to $(g_{\mathcal{H}})_{(u_0,t_0)}$. Take any $X,Y \in \mathcal{H}_{(u_0,t_0)}$. Let $\widetilde{X}$ be a section of $\mathcal{H}$ on a neighborhood of (u_0,t_0) with $\widetilde{X}_{(u_0,t_0)} = X$ and $(\nabla^{\mathcal{H}}X)_{(u_0,t_0)} = 0$. Similarly we define $\widetilde{Y}$ and $\widetilde{e}_i$. Let $W = X, Y$ or e_i. Then, from $(\nabla^{\mathcal{H}}_{e_i}\widetilde{W})_{(u_0,t_0)} = 0$, $(\nabla_{e_i}\widetilde{W})_{(u_0,t_0)} = \mathcal{A}_{e_i}W$ and the skew-symmetricness of $\mathcal{A}|_{\mathcal{H}\times\mathcal{H}}$, we can show

$$\begin{aligned}(\triangle_{\mathcal{H}}S)_{\mathcal{H}}(X,Y) &= \sum_{i=1}^{n}(\nabla_{e_i}\nabla_{e_i}S)(X,Y)\\ = \sum_{i=1}^{n}(\nabla^{\mathcal{H}}_{e_i}\nabla^{\mathcal{H}}_{e_i}S_{\mathcal{H}})(X,Y) &- 2\sum_{i=1}^{n}((\nabla_{e_i}S)(\mathcal{A}_{e_i}X,Y) + (\nabla_{e_i}S)(\mathcal{A}_{e_i}Y,X))\\ &- \sum_{i=1}^{n}(S(\mathcal{A}_{e_i}(\mathcal{A}_{e_i}X),Y) + S(\mathcal{A}_{e_i}(\mathcal{A}_{e_i}Y),X))\\ &- \sum_{i=1}^{n}(S((\nabla_{e_i}\mathcal{A})_{e_i}X,Y) + S((\nabla_{e_i}\mathcal{A})_{e_i}Y,X))\\ &-2\sum_{i=1}^{n}S(\mathcal{A}_{e_i}X,\mathcal{A}_{e_i}Y),\end{aligned}$$

which is equal to the right-hand side of the relation in the statement. This completes the proof. q.e.d.

Also we have the following Simons-type identity.

Lemma 3.5. *We have*

$$\triangle_{\mathcal{H}}h = \nabla d||H|| + ||H||(A^2)_{\sharp} - (\mathrm{Tr}\,(A^2)_{\mathcal{H}})h,$$

where $(A^2)_{\sharp}$ is the element of $\Gamma(\pi_M^*T^{(0,2)}M)$ defined by $(A^2)_{\sharp}(X,Y) := g(A^2X,Y)$ $(X,Y \in \pi_M^*TM)$.

Proof. Take $X, Y, Z, W \in \pi_M^*(TM)$. Since the ambient space V is flat, it follows from the Ricci's identity, the Gauss equation and the Codazzi equation that

$$\begin{aligned}
&(\nabla_X\nabla_Y h)(Z,W) - (\nabla_Z\nabla_W h)(X,Y)\\
&= (\nabla_X\nabla_Z h)(Y,W) - (\nabla_Z\nabla_X h)(Y,W)\\
&= h(X,Y)h(AZ,W) - h(Z,Y)h(AX,W)\\
&\quad + h(X,W)h(AZ,Y) - h(Z,W)h(AX,Y).
\end{aligned}$$

By using this relation, we obtain the desired relation. q.e.d.

Note. In the sequel, we omit the notation F_* for simplicity.

Define a section $\mathcal{R}$ of $\pi_M^*(\mathcal{H}^{(0,2)})$ by

$$\begin{aligned}
\mathcal{R}(X,Y) := {}& \mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} h(\mathcal{A}_\bullet(\mathcal{A}_\bullet X), Y) + \mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} h(\mathcal{A}_\bullet(\mathcal{A}_\bullet Y), X)\\
&+ \mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} h((\nabla_\bullet\mathcal{A})_\bullet X, Y) + \mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} h((\nabla_\bullet\mathcal{A})_\bullet Y, X)\\
&+ 2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} (\nabla_\bullet h)(\mathcal{A}_\bullet X, Y) + 2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} (\nabla_\bullet h)(\mathcal{A}_\bullet Y, X)\\
&+ 2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} h(\mathcal{A}_\bullet X, \mathcal{A}_\bullet Y) \qquad (X, Y \in \mathcal{H}).
\end{aligned}$$

By using Lemmas 3.3, 3.4 and 3.5, we can show the following evolution equation.

Theorem 3.6. *The second fundamental forms* $(h_{\mathcal{H}})_t$ *satisfies the following evolution equation:*

$$\begin{aligned}
\frac{\partial h_{\mathcal{H}}}{\partial t}(X,Y) = {}& (\bar{\triangle}_{\mathcal{H}} h_{\mathcal{H}})(X,Y) - 2||H||((A_{\mathcal{H}})^2)_\sharp(X,Y) - 2||H||((\mathcal{A}^\phi_\xi)^2)_\sharp(X,Y)\\
&+ \mathrm{Tr}\left((A_{\mathcal{H}})^2 - ((\mathcal{A}^\phi_\xi)^2)_{\mathcal{H}}\right) h_{\mathcal{H}}(X,Y) - \mathcal{R}(X,Y)
\end{aligned}$$

for $X, Y \in \mathcal{H}$.

Proof. Take $X, Y \in \mathcal{H}_{(u,t)}$. Easily we have

$$AX = A_{\mathcal{H}}X + \mathcal{A}^\phi_\xi X, \tag{3.1}$$

and

$$(A^2)_{\mathcal{H}}X = (A_{\mathcal{H}})^2X - (\mathcal{A}^\phi_\xi)^2X, \tag{3.2}$$

where we use

$$\left(\widetilde{\nabla}_W\xi\right)_{\widetilde{\mathcal{H}}} = \left(\widetilde{\nabla}_\xi W + [W,\xi]\right)_{\widetilde{\mathcal{H}}} = \left(\widetilde{\nabla}_\xi W\right)_{\widetilde{\mathcal{H}}} = \mathcal{A}_\xi W$$

for $W \in \Gamma(\widetilde{\mathcal{V}})$ because of $[W, \xi] \in \Gamma(\widetilde{\mathcal{V}})$. Also, since $\left[\frac{\partial}{\partial t}, \bar{X}_{\mathcal{H}}\right] \in \mathcal{V}$, we have

$$\left[\frac{\partial}{\partial t}, \bar{X}_{\mathcal{H}}\right] = 2||H||\mathcal{A}^{\phi}_{\xi}\bar{X}_{\mathcal{H}}. \tag{3.3}$$

From Lemma 3.3, (3.1), (3.2) and (3.3), we have

$$\begin{aligned}
&\frac{\partial h_{\mathcal{H}}}{\partial t}(X,Y) = \frac{\partial}{\partial t}(h_{\mathcal{H}}(\bar{X},\bar{Y})) = \frac{\partial}{\partial t}(h(\bar{X}_{\mathcal{H}},\bar{Y}_{\mathcal{H}})) = \frac{\partial}{\partial t}\langle \xi, \bar{X}_{\mathcal{H}}(\bar{Y}_{\mathcal{H}}F)\rangle \\
&= \langle \frac{\partial \xi}{\partial t}, \bar{X}_{\mathcal{H}}(\bar{Y}_{\mathcal{H}}F)\rangle + \langle \xi, \frac{\partial}{\partial t}\left(\bar{X}_{\mathcal{H}}(\bar{Y}_{\mathcal{H}}F)\right)\rangle \\
&= -\langle F_*(\mathrm{grad}_g||H||), \widetilde{\nabla}_X F_*\bar{Y}_{\mathcal{H}}\rangle + \langle \xi, X\left(\bar{Y}_{\mathcal{H}}\left(\frac{\partial F}{\partial t}\right)\right)\rangle \\
&\quad +\langle \xi, X([\frac{\partial}{\partial t}, \bar{Y}_{\mathcal{H}}]F)\rangle + \langle \xi, [\frac{\partial}{\partial t}, \bar{X}_{\mathcal{H}}](\bar{Y}_{\mathcal{H}}F)\rangle \\
&= -g(\mathrm{grad}_g||H||, \nabla_X\bar{Y}_{\mathcal{H}}) + X(\bar{Y}_{\mathcal{H}}||H||) - ||H||\langle \xi, \widetilde{\nabla}_X F_*(A(\bar{Y}_{\mathcal{H}})\rangle \\
&\quad +\langle \xi, \widetilde{\nabla}_X F_*([\frac{\partial}{\partial t}, \bar{Y}_{\mathcal{H}}])\rangle + \langle \xi, \widetilde{\nabla}_{[\frac{\partial}{\partial t}, \bar{X}_{\mathcal{H}}]}F_*\bar{Y}_{\mathcal{H}}\rangle \\
&= (\nabla d||H||)(X,Y) - ||H||h_{\mathcal{H}}(X, A_{\mathcal{H}}Y) + ||H||h(X, \mathcal{A}^{\phi}_{\xi}Y) + 2||H||h(\mathcal{A}^{\phi}_{\xi}X, Y) \\
&= (\nabla d||H||)(X,Y) - ||H||g_{\mathcal{H}}((A_{\mathcal{H}})^2X, Y) - 3||H||g((\mathcal{A}^{\phi}_{\xi})^2X, Y)
\end{aligned}$$

From this relation and the Simons-type identity in Lemma 3.5, we have

$$\begin{aligned}
\frac{\partial h_{\mathcal{H}}}{\partial t} &= \triangle_{\mathcal{H}}h - 2||H||((A_{\mathcal{H}})^2)_\sharp - 2||H||((\mathcal{A}^{\phi}_{\xi})^2)_\sharp \\
&\quad +\mathrm{Tr}\left((A_{\mathcal{H}})^2 - ((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}}\right)h_{\mathcal{H}}.
\end{aligned} \tag{3.4}$$

Substituting the relation in Lemma 3.4 into (3.4), we obtain the desired relation. q.e.d.

By using Lemma 3.2, we can show the following relation.

Lemma 3.7. *Let X and Y be local sections of $\mathcal{H}$ such that $g(X,Y)$ is constant. Then we have* $g(\nabla_{\frac{\partial}{\partial t}}X, Y) + g(X, \nabla_{\frac{\partial}{\partial t}}Y) = 2||H||h(X,Y)$.

Proof. From Lemma 3.2, we have

$$\begin{aligned}
\frac{\partial}{\partial t}g(X,Y) &= \frac{\partial g}{\partial t}(X,Y) + g(\nabla_{\frac{\partial}{\partial t}}X, Y) + g(X, \nabla_{\frac{\partial}{\partial t}}Y) \\
&= -2||H||h(X,Y) + g(\nabla_{\frac{\partial}{\partial t}}X, Y) + g(X, \nabla_{\frac{\partial}{\partial t}}Y).
\end{aligned}$$

On the other hand, we have $\frac{\partial}{\partial t}g(X,Y) = 0$. Therefore we obtain the desired relation. q.e.d.

Next we prepare the following lemma for $\mathcal{R}$.

Lemma 3.8. *For $X, Y \in \mathcal{H}$, we have*

$$
\begin{aligned}
\mathcal{R}(X,Y) = {} & 2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\Big(\langle(\mathcal{A}^{\phi}_{\bullet}X, \mathcal{A}^{\phi}_{\bullet}(A_{\mathcal{H}}Y))\rangle + \langle(\mathcal{A}^{\phi}_{\bullet}Y, \mathcal{A}^{\phi}_{\bullet}(A_{\mathcal{H}}X))\rangle\Big) \\
& +2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\Big(\langle(\mathcal{A}^{\phi}_{\bullet}X, \mathcal{A}^{\phi}_{Y}(A_{\mathcal{H}}\bullet))\rangle + \langle(\mathcal{A}^{\phi}_{\bullet}Y, \mathcal{A}^{\phi}_{X}(A_{\mathcal{H}}\bullet))\rangle\Big) \\
& +2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\Big(\langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_{\xi}Y, \mathcal{A}^{\phi}_{\bullet}X\rangle + \langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_{\xi}X, \mathcal{A}^{\phi}_{\bullet}Y\rangle\Big) \\
& +\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\Big(\langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_{\bullet}X, \mathcal{A}^{\phi}_{\xi}Y\rangle + \langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_{\bullet}Y, \mathcal{A}^{\phi}_{\xi}X\rangle\Big) \\
& +2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{T}^{\phi}_{\mathcal{A}^{\phi}_{\bullet}X}\xi, \mathcal{A}^{\phi}_{\bullet}Y\rangle,
\end{aligned}
\tag{3.5}
$$

where we omit F_*.

Proof. Take $e, X, Y \in \mathcal{H}$. Easily we have

$$
\begin{aligned}
(\nabla_e h)(\mathcal{A}_e X, Y) = {} & e(\langle\mathcal{A}^{\phi}_{e}X, \mathcal{A}^{\phi}_{\xi}Y\rangle) - h(\mathcal{A}_e(\mathcal{A}_e X), Y) \\
& -h((\nabla_e\mathcal{A})_e X, Y) - h(\mathcal{A}_e X, \mathcal{A}_e Y).
\end{aligned}
\tag{3.6}
$$

On the other hand, by simple calculation, we have $((\widetilde{\nabla}_e\mathcal{A}^{\phi})_X\xi)_{\widetilde{\mathcal{V}}} = -((\widetilde{\nabla}_e\mathcal{A}^{\phi})_{\xi}X)_{\widetilde{\mathcal{V}}}$. By using this relation, we can show

$$
e(\langle\mathcal{A}^{\phi}_{e}X, \mathcal{A}^{\phi}_{\xi}Y\rangle) = \langle(\widetilde{\nabla}_e\mathcal{A}^{\phi})_e X, \mathcal{A}^{\phi}_{\xi}Y) + \langle(\widetilde{\nabla}_e\mathcal{A}^{\phi})_{\xi}Y, \mathcal{A}^{\phi}_{e}X\rangle + h(\mathcal{A}_Y\mathcal{A}_X e, e\rangle.
\tag{3.7}
$$

Also, by simple calculations, we have

$$
\begin{aligned}
h(\mathcal{A}_e(\mathcal{A}_e X), Y) &= -\langle\mathcal{A}^{\phi}_{e}X, \mathcal{A}^{\phi}_{e}(A_{\mathcal{H}}Y)\rangle \\
h(\mathcal{A}_Y(\mathcal{A}_X e), e) &= \langle\mathcal{A}^{\phi}_{e}X, \mathcal{A}^{\phi}_{Y}(A_{\mathcal{H}}e)\rangle, \\
h((\nabla_e\mathcal{A})_e X, Y) &= \langle(\widetilde{\nabla}_e\mathcal{A}^{\phi})_e X, \mathcal{A}^{\phi}_{\xi}Y\rangle, \\
h(\mathcal{A}_e X, \mathcal{A}_e Y) &= -\langle{\mathcal{T}^{\phi}}_{\mathcal{A}^{\phi}_{e}X}\xi, \mathcal{A}^{\phi}_{e}Y\rangle.
\end{aligned}
\tag{3.8}
$$

From (3.6), (3.7) and the relations in (3.8), we have the desired relation. q.e.d.

Also, we prepare the following lemma.

Lemma 3.9. *For $X, Y, Z \in \mathcal{H}$, we have*

$$
2\langle\mathcal{T}^{\phi}_{\mathcal{A}^{\phi}_{X}Y}\xi, \mathcal{A}^{\phi}_{X}Z\rangle = -\langle\mathcal{A}^{\phi}_{X}Z, (\widetilde{\nabla}_X\mathcal{A}^{\phi})_{\xi}Y\rangle + \langle\mathcal{A}^{\phi}_{X}Z, (\widetilde{\nabla}_Y\mathcal{A}^{\phi})_{\xi}X\rangle.
$$

Proof. Fix $(u_0,t_0)\in M\times[0,T)$. Let $\widetilde{X}$ be an element of $\Gamma(\mathcal{H})$ satisfying $\widetilde{X}_{(u_0,t_0)}=X$ and $(\widetilde{\nabla}^{\mathcal{H}}\widetilde{X})_{(u_0,t_0)}=0$. Let $\widetilde{Y}$ and $\widetilde{Z}$ be similar elements of $\Gamma(\mathcal{H})$ for Y and Z, respectively. At (u_0,t_0), we have

$$\begin{aligned}\langle\mathcal{A}^{\phi}_XZ,\mathcal{A}^{\phi}_Y(A_{\mathcal{H}}X)\rangle&=-\langle\mathcal{A}^{\phi}_XZ,\widetilde{\nabla}_Y(\widetilde{\nabla}_X\xi)-\widetilde{\nabla}_Y(\mathcal{A}^{\phi}_X\xi)\rangle\\&=-\langle\mathcal{A}^{\phi}_XZ,\widetilde{\nabla}_X(\widetilde{\nabla}_Y\xi)+\widetilde{\nabla}_{[Y,X]}\xi-\widetilde{\nabla}_Y(\mathcal{A}^{\phi}_X\xi)\rangle\\&=\langle\mathcal{A}^{\phi}_XZ,\widetilde{\nabla}_X(AY)\rangle-2\langle\mathcal{A}^{\phi}_XZ,\widetilde{\nabla}_{\mathcal{A}^{\phi}_YX}\xi\rangle\\&\qquad+\langle\mathcal{A}^{\phi}_XZ,(\widetilde{\nabla}_Y\mathcal{A}^{\phi})_X\xi\rangle-\langle\mathcal{A}^{\phi}_XZ,\mathcal{A}^{\phi}_X(AY)\rangle\\&=\langle\mathcal{A}^{\phi}_XZ,\mathcal{A}^{\phi}_Y(A_{\mathcal{H}}X)\rangle-\langle\mathcal{A}^{\phi}_XZ,(\widetilde{\nabla}_X\mathcal{A}^{\phi})_Y\xi\rangle\\&\qquad-2\langle\mathcal{A}^{\phi}_XZ,\mathcal{T}^{\phi}_{\mathcal{A}^{\phi}_YX}\xi\rangle+\langle\mathcal{A}^{\phi}_XZ,(\widetilde{\nabla}_Y\mathcal{A}^{\phi})_X\xi\rangle,\end{aligned}\tag{3.9}$$

where we use $(\widetilde{\nabla}^{\mathcal{H}}\widetilde{X})_{(u_0,t_0)}=(\widetilde{\nabla}^{\mathcal{H}}\widetilde{Y})_{(u_0,t_0)}=(\widetilde{\nabla}^{\mathcal{H}}\widetilde{Z})_{(u_0,t_0)}=0$. Also we have

$$\langle\mathcal{A}^{\phi}_XZ,(\widetilde{\nabla}_X\mathcal{A}^{\phi})_Y\xi\rangle=-\langle\mathcal{A}^{\phi}_XZ,(\widetilde{\nabla}_X\mathcal{A}^{\phi})_\xi Y\rangle$$

and

$$\langle\mathcal{A}^{\phi}_XZ,(\widetilde{\nabla}_Y\mathcal{A}^{\phi})_X\xi\rangle=-\langle\mathcal{A}^{\phi}_XZ,(\widetilde{\nabla}_Y\mathcal{A}^{\phi})_\xi X\rangle.$$

Form (3.9) and these relations, we obtain the desired relation. q.e.d.

Lemma 3.10. *For $X\in\mathcal{H}$, we have*

$$\begin{aligned}\mathcal{R}(X,X)=&\,4\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{A}^{\phi}_{\bullet}X,\mathcal{A}^{\phi}_{\bullet}(A_{\mathcal{H}}X)\rangle+4\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{A}^{\phi}_{\bullet}X,\mathcal{A}^{\phi}_X(A_{\mathcal{H}}\bullet)\rangle\\&+3\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_\xi X,\mathcal{A}^{\phi}_{\bullet}X\rangle+2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_{\bullet}X,\mathcal{A}^{\phi}_\xi X\rangle\\&+\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{A}^{\phi}_{\bullet}X,(\widetilde{\nabla}_X\mathcal{A}^{\phi})_\xi\bullet\rangle\end{aligned}$$

and hence

$$\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\mathcal{R}(\bullet,\bullet)=0.$$

Proof. These relations follow from the relations in Lemmas 3.8 and 3.9 directly. q.e.d.

By using Theorem 3.6 and Lemmas 3.7 and 3.10, we can show the following evolution equation.

Corollary 3.11. *The norms $||H_t||$ of H_t satisfies the following evolution equation:*

$$\frac{\partial||H||}{\partial t}=\triangle_{\mathcal{H}}||H||+||H||\mathrm{Tr}(A_{\mathcal{H}})^2-3||H||\mathrm{Tr}((\mathcal{A}^{\phi}_\xi)^2)_{\mathcal{H}}.$$

Proof. Fix $(u_0, t_0) \in M \times [0, T)$. Take a local orthonormal frame field $\{e_1, \cdots, e_n\}$ of $\mathcal{H}$ over a neighborhood U of (u_0, t_0) consisting of the eigenvectors of $A_{\mathcal{H}}$, where the orthonomality means one for g. Since the fibres of ϕ are regularized minimal Fredholm submanifolds, we have $||H|| = \sum_{i=1}^{n} h(e_i, e_i)$ on U. Clearly we have

$$\frac{\partial ||H||}{\partial t} = \sum_{i=1}^{n} \left(\frac{\partial h_{\mathcal{H}}}{\partial t}(e_i, e_i) + 2h_{\mathcal{H}}(\nabla_{\frac{\partial}{\partial t}} e_i, e_i) \right). \tag{3.10}$$

On the other hand, it follows from Theorem 3.6 that

$$\sum_{i=1}^{n} \frac{\partial h_{\mathcal{H}}}{\partial t}(e_i, e_i) = \triangle_{\mathcal{H}} ||H|| - ||H|| \mathrm{Tr}(A_{\mathcal{H}})^2 - 3||H|| \mathrm{Tr}((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}}, \tag{3.11}$$

where we use $\sum_{i=1}^{n} (\bar{\triangle}_{\mathcal{H}} h_{\mathcal{H}})(e_i, e_i) = \triangle_{\mathcal{H}} ||H||$ and $\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} \mathcal{R}(\bullet, \bullet) = 0$ (by Lemma 3.10). Since each e_i is an eigenvector of $A_{\mathcal{H}}$, we have $h(e_i, e_j) = 0$ $(i \neq j)$. By using Lemma 3.7, we can show

$$\sum_{i=1}^{n} h_{\mathcal{H}}(\nabla_{\frac{\partial}{\partial t}} e_i, e_i) = \sum_{i=1}^{n} g(\nabla_{\frac{\partial}{\partial t}} e_i, e_i) h(e_i, e_i) = ||H|| \mathrm{Tr}(A_{\mathcal{H}})^2. \tag{3.12}$$

From (3.10), (3.11) and (3.12), we obtain the desired relation. q.e.d.

By using Lemmas 3.2, 3.6 and 3.7, we can show the following evolution equation.

Corollary 3.12. *The quantities* $\mathrm{Tr}(A_{\mathcal{H}})^2_t$ *satisfies the following evolution equation:*

$$\begin{aligned} \frac{\partial \mathrm{Tr}(A_{\mathcal{H}})^2}{\partial t} = & \triangle_{\mathcal{H}}(\mathrm{Tr}(A_{\mathcal{H}})^2) - 2\mathrm{Tr}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}(\nabla^{\mathcal{H}}_{\bullet} A_{\mathcal{H}} \circ \nabla^{\mathcal{H}}_{\bullet} A_{\mathcal{H}}) \\ & + 2\mathrm{Tr}((A_{\mathcal{H}})^2) \left(\mathrm{Tr}((A_{\mathcal{H}})^2) - \mathrm{Tr}((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}} \right) \\ & - 4||H|| \mathrm{Tr} \left(((\mathcal{A}^{\phi}_{\xi})^2) \circ A_{\mathcal{H}} \right) - 2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} \mathcal{R}(A_{\mathcal{H}} \bullet, \bullet). \end{aligned}$$

Proof. Fix $(u_0, t_0) \in M \times [0, T)$. Take a local orthonormal frame field $\{e_1, \cdots, e_n\}$ of $\mathcal{H}$ over a neighborhood U of (u_0, t_0) consisting of the eigenvectors of $A_{\mathcal{H}}$. From Lemma 3.2, we have

$$\begin{aligned} \frac{\partial h_{\mathcal{H}}}{\partial t}(X, Y) &= \frac{\partial g_{\mathcal{H}}}{\partial t}(A_{\mathcal{H}} X, Y) + g_{\mathcal{H}}(\frac{\partial A_{\mathcal{H}}}{\partial t}(X), Y) \\ &= -2||H|| h_{\mathcal{H}}(A_{\mathcal{H}} X, Y) + g_{\mathcal{H}}(\frac{\partial A_{\mathcal{H}}}{\partial t}(X), Y) \end{aligned} \tag{3.13}$$

for any $X, Y \in \pi_M^* TM$. Since $\{e_1, \cdots, e_n\}$ consists of the eigenvectors of $A_{\mathcal{H}}$, it follows from Lemma 3.7 that

$$g(\nabla_{\frac{\partial}{\partial t}} e_i, e_i) = ||H|| h(e_i, e_i). \tag{3.14}$$

From these relations, Lemmas 3.2 and 3.7, we have
(3.15)

$$\begin{aligned}
&\frac{\partial \mathrm{Tr}(A_{\mathcal{H}})^2}{\partial t} = \sum_{i=1}^{n} \frac{\partial}{\partial t}(h_{\mathcal{H}}(A_{\mathcal{H}} e_i, e_i)) \\
&= \sum_{i=1}^{n} \left(\frac{\partial h_{\mathcal{H}}}{\partial t}(A_{\mathcal{H}} e_i, e_i) + h_{\mathcal{H}}(\frac{\partial A_{\mathcal{H}}}{\partial t}(e_i), e_i) + 2h_{\mathcal{H}}(A_{\mathcal{H}} e_i, \nabla_{\frac{\partial}{\partial t}} e_i) \right) \\
&= \sum_{i=1}^{n} \left(\frac{\partial h_{\mathcal{H}}}{\partial t}(A_{\mathcal{H}} e_i, e_i) + g_{\mathcal{H}}((\frac{\partial A_{\mathcal{H}}}{\partial t}(e_i), A_{\mathcal{H}} e_i) + 2||H|| h(e_i, e_i) h_{\mathcal{H}}(A_{\mathcal{H}} e_i, e_i) \right) \\
&= \sum_{i=1}^{n} \left(2\frac{\partial h_{\mathcal{H}}}{\partial t}(A_{\mathcal{H}} e_i, e_i) + 2||H|| g((A_{\mathcal{H}})^3 e_i, e_i) + 2||H|| h(e_i, e_i) g((A_{\mathcal{H}})^2 e_i, e_i) \right) \\
&= \sum_{i=1}^{n} \left(2\frac{\partial h_{\mathcal{H}}}{\partial t}(A_{\mathcal{H}} e_i, e_i) + 4||H|| g((A_{\mathcal{H}})^3 e_i, e_i) \right).
\end{aligned}$$

Also we have

$$\sum_{i=1}^{n} (\bar{\triangle}_{\mathcal{H}} h_{\mathcal{H}})(A_{\mathcal{H}} e_i, e_i) = \frac{1}{2} \triangle_{\mathcal{H}} \mathrm{Tr}((A_{\mathcal{H}})^2) - \mathrm{Tr}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} \left(\nabla^{\mathcal{H}}_{\bullet} A_{\mathcal{H}} \circ \nabla^{\mathcal{H}}_{\bullet} A_{\mathcal{H}} \right). \tag{3.16}$$

From Theorem 3.6, (3.15) and (3.16), we obtain the desired relation. q.e.d.

By using Corollaries 3.8 and 3.9, we can show the following evolution equation.

Corollary 3.13. *The quantities* $\mathrm{Tr}\,(A_{\mathcal{H}})^2_t - \frac{||H_t||^2}{n}$ *satisfies the following evolution equation:*

$$\begin{aligned}
\frac{\partial (\mathrm{Tr}\,(A_{\mathcal{H}})^2 - \frac{||H||^2}{n})}{\partial t} = {}& \triangle_{\mathcal{H}} \left(\mathrm{Tr}(A_{\mathcal{H}})^2 - \frac{||H||^2}{n} \right) + \frac{2}{n} ||\mathrm{grad}||H||||^2 \\
& + 2\mathrm{Tr}(A_{\mathcal{H}})^2 \times \left(\mathrm{Tr}(A_{\mathcal{H}})^2 - \frac{||H||^2}{n} \right) \\
& - 2\mathrm{Tr}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} \left(\nabla^{\mathcal{H}} A_{\mathcal{H}} \circ \nabla^{\mathcal{H}} A_{\mathcal{H}} \right) \\
& - 2\mathrm{Tr}((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}} \times \left(\mathrm{Tr}(A_{\mathcal{H}})^2 - \frac{||H||^2}{n} \right) \\
& - 4||H|| \left(\mathrm{Tr} \left((\mathcal{A}^{\phi}_{\xi})^2 \circ \left(A_{\mathcal{H}} - \frac{||H||}{n} \mathrm{id} \right) \right) \right) \\
& - 2\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}} \mathcal{R} \left(\left(A_{\mathcal{H}} - \frac{||H||}{n} \mathrm{id} \right) \bullet, \bullet \right),
\end{aligned}$$

where $\mathrm{grad}||H||$ *is the gradient vector field of* $||H||$ *with respect to* g *and* $||\mathrm{grad}||H||||$ *is the norm of* $\mathrm{grad}||H||$ *with respect to* g*.*

Proof. This relation follows directly from Corollaries 3.8, 3.9 and $\triangle_{\mathcal{H}}||H||^2 = 2||H||\triangle_{\mathcal{H}}||H|| + 2\,||\mathrm{grad}||H||||^2$. q.e.d.

Remark 3.1. From the evolution equations in Corollary 3.11 $\sim$ 3.13, the evolution equations for the corresponding geometric quantities of $\bar{f}_t(: \bar{M} \hookrightarrow V/G)$ are derived, respectively. These derived evolution equations coincides with the evolution equations obtained by Huisken [Hu2], where we note that Huisken [Hu2] treat the mean curvature flow in a complete Riemannian manifold in general. Namely the discussion in this section give a new proof of the evolution equations in [Hu2] in the case where the ambient space is equal to V/G.

4 A maximum principle

Let M be a Hilbert manifold and g_t $(0 \leq t < T)$ a C^∞-family of Riemannian metrics on M and $G \curvearrowright M$ a free action on M such that G-action is isometric with respect to g_t and that the orbit space M/G is compact. Let $\mathcal{H}_t$ $(0 \leq t < T)$ be the horizontal distribution of the G-action. Let π_M be the natural projection of $M \times [0,T)$ onto M. For a tangent vector field X on M (or an open set U of M), we define a section $\bar{X}$ of π_M^*TM (or $\pi_M^*TM|_U$) by $\bar{X}_{(x,t)} := X_x$ $((x,t) \in M \times [0,T))$. Denote by ∇^t $(0 \leq t < T)$ the Riemannian connection of g_t. Define a connection ∇ of π_M^*TM by $(\nabla_{\bar{X}}\bar{Y})_{(x,t)} = (\nabla^t_X Y)_x$ and $\nabla_{\frac{\partial}{\partial t}}\bar{X} = \nabla_{\bar{X}}\frac{\partial}{\partial t} = 0$ for $X, Y \in \Gamma(TM)$. Define a subbundle $\mathcal{H}$ of π_M^*TM by $\mathcal{H}_{(x,t)} := (\mathcal{H}_t)_x$. Define a connection $\nabla^{\mathcal{H}}$ of $\mathcal{H}$ by $\nabla^{\mathcal{H}}_X Y = \mathrm{pr}_{\mathcal{H}}(\nabla_X Y)$ for any $X \in T(M \times [0,T))$ and any $Y \in \Gamma(\mathcal{H})$, where $\mathrm{pr}_{\mathcal{H}}$ is the orthogonal projection of π_M^*TM onto $\mathcal{H}$. Denote by $\Gamma(\pi_M^*T^{(r,s)}M)$ the space of all sections of $\pi_M^*(T^{(r,s)}M)$. Let $B \in \Gamma(\pi_M^*T^{(r_0,s_0)}M)$ We define maps $\psi_{B\otimes}$ and $\psi_{\otimes B}$ from $\Gamma(\pi_M^*T^{(r,s)}M)$ to $\Gamma(\pi_M^*T^{(r+r_0,s+s_0)}M)$ by

$$\psi_{B\otimes}(S) := B \otimes S, \quad \text{and} \quad \psi_{\otimes B}(S) := S \otimes B \quad (S \in \Gamma(\pi_M^*T^{(r,s)}M),$$

respectively. Also, we define a map $\psi_{\otimes^k}$ of $\Gamma(\pi_M^*T^{(r,s)}M)$ to $\Gamma(\pi_M^*T^{(kr,ks)}M)$ by

$$\psi_{\otimes^k}(S) := S \otimes \cdots \otimes S \ (k\text{−times}) \quad (S \in \Gamma(\pi_M^*T^{(r,s)}M).$$

Also, we define a map $\psi_{g_{\mathcal{H}},ij}$ $(i < j)$ from $\Gamma(\pi_M^* T^{(0,s)}M)$ (or $\Gamma(\pi_M^* T^{(1,s)}M)$) to $\Gamma(\pi_M^* T^{(0,s-2)}M)$ (or $\Gamma(\pi_M^* T^{(1,s-2)}M)$) by

$$\begin{aligned}&(\psi_{g_{\mathcal{H}},ij}(S))_{(x,t)}(X_1,\cdots,X_{s-2})\\ :=&\sum_{k=1}^{n} S_{(u,t)}(X_1,\cdots,X_{i-1},e_k,X_{i+1},\cdots,X_{j-1},e_k,X_{j+1},\cdots,X_{s-2})\end{aligned}$$

and define a map $\psi_{\mathcal{H},i}$ from $\Gamma(\pi_M^* T^{(1,s)}M)$ to $\Gamma(\pi_M^* T^{(0,s-1)}M)$ by

$$(\psi_{\mathcal{H},i}(S))_{(u,t)}(X_1,\cdots,X_{s-1}) := \mathrm{Tr} S_{(u,t)}(X_1,\cdots,X_{i-1},\bullet,X_i,\cdots,X_{s-1}),$$

where $X_i \in T_xM$ $(i = 1,\cdots,s-1)$ and $\{e_1,\cdots,e_n\}$ is an orthonormal base of $(\mathcal{H}_t)_x$ with respect to g_t. We call a map P from $\Gamma(\pi_M^* T^{(0,s)}M)$ to oneself given by the composition of the above maps of five type *a map of polynomial type.*

In this section, we prove the following maximum principle for a C^∞-family of $(0,2)$-tensor field on M.

Theorem 4.1. *Let $S \in \Gamma(\pi_M^*(T^{(0,2)}M))$ such that, for each $t \in [0,T)$, $S_t(:= S_{(\cdot,t)})$ is a G-invariant symmetric section of $T^{(0,2)}M$. Assume that S_t $(0 \le t < T)$ satisfy the following evolution equation:*

$$\frac{\partial S_{\mathcal{H}}}{\partial t} = \bar{\triangle}_{\mathcal{H}} S_{\mathcal{H}} + \nabla^{\mathcal{H}}_{X_0} S_{\mathcal{H}} + P(S)_{\mathcal{H}}, \tag{4.1}$$

where $X_0 \in \Gamma(TM)$ and P is a map of polynomial type from $\Gamma(\pi_M^*(T^{(0,2)}M))$ to oneself.

(i) *Assume that P satisfies the following condition:*

$$X \in \mathrm{Ker}(\hat{S}_{\mathcal{H}})_{(x,t)} \ \Rightarrow\ P(\hat{S})_{(x,t)}(X,X) \ge 0$$

for each $\hat{S} \in \Gamma(\pi_M^*(T^{(0,2)}M))$ and each $(x,t) \in M \times [0,T)$, Then, if $(S_{\mathcal{H}})_{(\cdot,0)} \ge 0$ (resp. > 0), then $(S_{\mathcal{H}})_{(\cdot,t)} \ge 0$ (resp. > 0) holds for all $t \in [0,T)$.

(ii) *Assume that P satisfies the following condition:*

$$X \in \mathrm{Ker}(\hat{S}_{\mathcal{H}})_{(x,t)} \ \Rightarrow\ P(\hat{S})_{(x,t)}(X,X) \le 0$$

for each $\hat{S} \in \Gamma(\pi_M^*(T^{(0,2)}M))$ and each $(x,t) \in M \times [0,T)$. Then, if $(S_{\mathcal{H}})_{(\cdot,0)} \le 0$ (resp. < 0), then $(S_{\mathcal{H}})_{(\cdot,t)} \le 0$ (resp. < 0) holds for all $t \in [0,T)$.

Proof. We shall show that, if $(S_{\mathcal{H}})_{(\cdot,0)} \ge 0$, then $(S_{\mathcal{H}})_{(\cdot,t)} \ge 0$ holds for all $t \in (0,T)$ in the statement (i). Note that the remained statement in the statement (i) and the statement (ii) also are shown similarly. For positive numbers ε and δ, we define $S_{\varepsilon,\delta}$ by $(S_{\varepsilon,\delta})_{(x,t)} := S_{(x,t)} + \varepsilon(\delta + t)g_{(x,t)}$.

(Step I) In this step, we show the following statement:

$$(*)\quad \exists\,\delta > 0 \text{ s.t. } "((S_{\varepsilon,\delta})_{\mathcal{H}})_{(x,t)} > 0\ (\forall\,(x,t) \in M \times [0,\delta), \forall\,\varepsilon > 0)".$$

Suppose that such a positive number δ does not exists. Fix a sufficiently small positive number δ. For some $\varepsilon_0 > 0$, there exists $(x_0,t_0) \in M \times [0,\delta)$ such that $((S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)} > 0$ does not hold. Here we take t_0 as smally as possible. We have $\mathrm{Ker}((S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)} \neq \{0\}$ and $((S_{\varepsilon_0,\delta})_t)_{\mathcal{H}_t} \geq 0$ $(\forall\,t \in [0,t_0])$. Take $v_1 \in \mathrm{Ker}((S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)})$ with $g(v_1,v_1) = 1$. From the assumption for P, we have

$$P((S_{\varepsilon_0,\delta})_{(x_0,t_0)})(v_1,v_1) \geq 0. \tag{4.2}$$

Since P is of polynomial type, M/G is compact and S_t is G-invariant, there exists a positive constant $C_{\delta,t}$ (depending on only $||S_{\mathcal{H}}||$ and $||(S_{\varepsilon_0,\delta})_{\mathcal{H}}||$) such that

$$||(P(S_{\varepsilon_0,\delta}))_{\mathcal{H}} - (P(S))_{\mathcal{H}}|| \leq C_{\delta,t}||(S_{\varepsilon_0,\delta})_{\mathcal{H}} - S_{\mathcal{H}}|| \tag{4.3}$$

on $M \times \{t\}$ for each $t \in [0,T)$, where $||\cdot||$ is the pointwise norm of a tensor field $(\cdot)$. We take $C_{\delta,t}$ as smally as possible. Since P is of polynomial type, $\lim_{\delta\to+0} C_{\delta,t}$ exists and $\lim_{\delta\to+0} C_{\delta,t} > 0$. Denote by C_t this limit. Fix $T_1 \in (t_0,T)$. Set

$$C_\delta := \max\left\{ \max_{0\leq t\leq T_1} C_{\delta,t},\quad \max_{\substack{(x,t)\in M\times[0,T_1]\\ v\in TM \text{ s.t. } g_t(v,v)=1}} \left|\left(\frac{\partial g_{\mathcal{H}}}{\partial t}\right)_{(x,t)}(v,v)\right| \right\}$$

and

$$C := \max\left\{ \max_{0\leq t\leq T_1} C_t,\quad \max_{\substack{(x,t)\in M\times[0,T_1]\\ v\in TM \text{ s.t. } g_t(v,v)=1}} \left|\left(\frac{\partial g_{\mathcal{H}}}{\partial t}\right)_{(x,t)}(v,v)\right| \right\}.$$

Since C is independent of the choice of δ, we may assume that $C\delta < \frac{1}{4}$ by replacing δ to a smaller positive number if necessary. Furthermore, since $\delta \mapsto C_\delta$ is upper semi-continuous, we may assume that $C_\delta\delta < \frac{1}{4}$ by replacing δ to a smaller positive number if necessary. From (4.2) and (4.3), we have

$$P(S)_{(x_0,t_0)}(v_1,v_1) \geq -2C_\delta\varepsilon_0\delta. \tag{4.4}$$

Let X_1 be a section of $\mathcal{H}$ on a normal neighborhood U of (x_0, t_0) in $M \times [0, T)$ such that $(X_1)_{(x_0,t_0)} = v_1$ and that $\nabla^{\mathcal{H}} X_1 = 0$ at (x_0, t_0). Define a function ρ on U by $\rho(x, t) := (S_{\varepsilon_0,\delta})_{(x,t)}((\bar{X}_1)_{(x,t)}, (\bar{X}_1)_{(x,t)})$ $((x, t) \in U)$. Since we take (x_0, t_0) and v_1 as above, we have $(\frac{\partial \rho}{\partial t})_{(x_0,t_0)} \leq 0$ (see Fig. 1). Also, we have

$$(\frac{\partial \rho}{\partial t})_{(x_0,t_0)} = (\frac{\partial S_{\mathcal{H}}}{\partial t})_{(x_0,t_0)}(v_1, v_1) + \varepsilon_0(\delta + t_0)(\frac{\partial g_{\mathcal{H}}}{\partial t})_{(x_0,t_0)}(v_1, v_1) + \varepsilon_0.$$

Hence we have

$$(\frac{\partial S_{\mathcal{H}}}{\partial t})_{(x_0,t_0)}(v_1, v_1) \leq -\varepsilon_0(\delta + t_0)(\frac{\partial g_{\mathcal{H}}}{\partial t})_{(x_0,t_0)}(v_1, v_1) - \varepsilon_0. \tag{4.5}$$

Take $w \in T_{x_0}(M \times \{t_0\})$. Clearly we have $d\rho_{(x_0,t_0)}(w) = 0$. Also we have $d\rho_{(x_0,t_0)}(w) = (\nabla^{\mathcal{H}}_w (S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)}(v_1, v_1)$. Hence we have

$$(\nabla^{\mathcal{H}}_w (S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)}(v_1, v_1) = 0. \tag{4.6}$$

Clearly we have $(\triangle^{t_0} \rho_{t_0})_{x_0} \geq 0$, where $\triangle^{t_0}$ is the Laplacian operator with respect to g_{t_0}. Also, we have $(\triangle^{t_0} \rho_{t_0})_{x_0} = (\bar{\triangle}_{\mathcal{H}}(S_{\varepsilon_0,\delta})_{\mathcal{H}})(v_1, v_1)$. Hence we have

$$(\bar{\triangle}_{\mathcal{H}}(S_{\varepsilon_0,\delta})_{\mathcal{H}})(v_1, v_1) \geq 0. \tag{4.7}$$

From the evolution equation in the statement, (4.5), (4.6) and (4.7), we have

$$\begin{aligned} P(S)_{(x_0,t_0)}(v_1, v_1) &\leq -\varepsilon_0 + \varepsilon_0(\delta + t_0) \left| (\frac{\partial g_{\mathcal{H}}}{\partial t})_{(x_0,t_0)}(v_1, v_1) \right| \\ &\leq -\varepsilon_0 + 2\varepsilon_0 C_\delta \delta. \end{aligned} \tag{4.8}$$

From (4.4) and (4.8), we have $C_\delta \delta \geq \frac{1}{4}$. This contradicts $C_\delta \delta < \frac{1}{4}$. Therefore the statement $(*)$ is true.

(Step II) Let δ be a positive number as in the statement $(*)$. Then, for any $(x, t) \in M \times [0, \delta)$ and any $\varepsilon > 0$, we have $((S_{\varepsilon,\delta})_{\mathcal{H}})_{(x,t)} > 0$. Hence we have $\lim_{\varepsilon \to +0} ((S_{\varepsilon,\delta})_{\mathcal{H}})_{(x,t)} = (S_{\mathcal{H}})_{(x,t)} \geq 0$ for any $(x, t) \in M \times [0, \delta)$. Set

$$T_1 := \sup\{t_1 \,|\, (S_{\mathcal{H}})_{(x,t)} \geq 0 \ (\forall\, t \in [0, t_1], \forall\, x \in M)\}.$$

Suppose that $T_1 < T$. Then, by the similar discussion for $(S_{\mathcal{H}})_{(\cdot,T_1)}$ instead of $(S_{\mathcal{H}})_{(\cdot,0)}$, we can show that $(S_{\mathcal{H}})_{(x,t)} \geq 0$ for any $t \in [T_1, T_1 + \delta']$ and any $x \in M$, where δ' is some positive number. This contradicts the definition of T_1. Therefore we have $T_1 = T$. Thus we obtain $(S_{\mathcal{H}})_{(\cdot,t)} \geq 0$ for any $t \in [0, T)$. This completes the proof. q.e.d.



Fig. 1.

Remark 4.1. The constant C_δ in this proof corresponds to the constant C in the proof of Theorem 9.1 in [Ha].

Similarly we obtain the following maximal principle for a C^∞-family of G-invariant functions on M.

Theorem 4.2. *Let ρ be a C^∞-function over $M \times [0,T)$ such that, for each $t \in [0,T)$, $\rho_t (:= \rho(\cdot,t))$ is a G-invariant function on M. Assume that ρ_t $(0 \le t < T)$ satisfy the following evolution equation:*

$$\frac{\partial \rho}{\partial t} = \triangle_{\mathcal{H}} \rho + d\rho(\bar{X}_0) + P(\rho),$$

where $X_0 \in \Gamma(TM)$ and P is a map of polynomial type from $C^\infty(M \times [0,T))$ to oneself.

(i) *Assume that P satisfies the following condition:*

$$\hat{\rho}_{(x_0,t_0)} = 0 \;\Rightarrow\; P(\hat{\rho})_{(x_0,t_0)} \ge 0$$

for each $\hat{\rho} \in C^\infty(M \times [0,T))$. Then, if $\rho_0 \ge 0$ (resp. > 0), then $\rho_t \ge 0$ (resp. > 0) holds for all $t \in [0,T)$.

(ii) *Assume that P satisfies the following condition:*

$$\hat\rho_{(x_0,t_0)} = 0 \ \Rightarrow \ P(\hat\rho)_{(x_0,t_0)} \le 0$$

for each $\hat\rho \in C^\infty(M \times [0,T))$. Then, if $\rho_0 \le 0$ (resp. < 0), then $\rho_t \le 0$ (resp. < 0) holds for all $t \in [0,T)$.

5 Strictly horizontally convexity preservability theorem

Let $G \curvearrowright V$ be an isometric free action with minimal Fredholm orbit of a Hilbert Lie group G on a Hilbert space V equipped with an inner product $\langle\ ,\ \rangle$ and $\phi : V \to V/G$ the orbit map. Denote by $\widetilde\nabla$ the Riemannian connection of V. Assume that V/G is compact. Set $n+1 := \dim V/G$. Let $M(\subset V)$ be a G-invariant hypersurface in V such that $\phi(M)$ is compact. Let f be an inclusion map of M into V and f_t $(0 \le t < T)$ the regularized mean curvature flow starting from f. We use the notations in Section 3. In the sequel, we omit the notation f_{t*} for simplicity. Set

$$L := \max_{(X_1,\cdots,X_5)\in\widetilde{\mathcal{H}}_1^5} |\langle \mathcal{A}^\phi_{X_1}((\widetilde\nabla_{X_2}\mathcal{A}^\phi)_{X_3}X_4),\ X_5\rangle|,$$

where $\widetilde{\mathcal{H}}_1 := \{X \in \widetilde{\mathcal{H}} \mid ||X|| = 1\}$. Note that L exists because of the compactness of V/G. In this section, we prove the following strictly horizontally convexity-preservability theorem by using results stated in Section 3 and the discussion in the proof of Theorem 4.1.

Theorem 5.1. *If M satisfies $||H_0||^2(h_{\mathcal{H}})_{(\cdot,0)} > 2n^2L(g_{\mathcal{H}})_{(\cdot,0)}$, then $T < \infty$ holds and $||H_t||^2(h_{\mathcal{H}})_{(\cdot,t)} > 2n^2L(g_{\mathcal{H}})_{(\cdot,t)}$ holds for all $t \in [0,T)$.*

Proof. Since $\mathcal{A}^\phi_\xi$ is skew-symmetric, we have

$$\mathrm{Tr}((\mathcal{A}^\phi_\xi)^2)_{\mathcal{H}} \le 0. \tag{5.1}$$

From Corollary 3.11, $\mathrm{Tr}(A_{\mathcal{H}})^2 \ge \frac{||H||^2}{n}$ and (5.1), we have

$$\frac{\partial ||H||}{\partial t} \ge \triangle_{\mathcal{H}} ||H|| + \frac{||H||^3}{n}. \tag{5.2}$$

Define a function ρ over $[0,T)$ by $\rho(t) := \min ||H_t||$. Form (5.2), we have $\frac{d\rho}{dt} \ge \frac{1}{n}\rho^3$. Also we have $\rho(0) > 0$ by the assumption. Hence we obtain $T \le \frac{n}{2\rho(0)^2}$.

Set $S := \dfrac{1}{||H||}h - \dfrac{2n^2L}{||H||^3}g$ and $S_{\varepsilon,\delta} := S + \varepsilon(\delta + t)g$, where ε and δ are positive constants. We shall show the following statement:

(*) $\exists\,\delta > 0$ s.t. $(S_{\varepsilon,\delta})_{(x,t)} > 0$ $(\forall\,(x,t) \in M \times [0,\delta),\ \forall\,\varepsilon > 0)$.

Take $X, Y \in \mathcal{H}$. By using Lemma 3.2, Theorem 3.6, Corollary 3.8 and Lemma 3.10, we can show

$$\begin{aligned}
&\frac{\partial (S_{\varepsilon,\delta})_{\mathcal{H}}}{\partial t}(X,Y)\\
&= \frac{1}{||H||}(\bar{\triangle}_{\mathcal{H}} h_{\mathcal{H}})(X,Y) - 2((A_{\mathcal{H}})^2)_\sharp(X,Y) - 2((\mathcal{A}^\phi_\xi)^2)_\sharp(X,Y)\\
&\quad -\frac{1}{||H||^2}\left(\bar{\triangle}_{\mathcal{H}}||H|| - 2||H||\mathrm{Tr}((\mathcal{A}^\phi_\xi)^2)_{\mathcal{H}} - 4n^2L\right) h_{\mathcal{H}}(X,Y)\\
&\quad -\frac{1}{||H||}\mathcal{R}(X,Y) + \varepsilon g_{\mathcal{H}}(X,Y) - 2\varepsilon(\delta+t)||H||h_{\mathcal{H}}(X,Y)\\
&\quad +\frac{3n^2L}{||H||^4}\left(\bar{\triangle}_{\mathcal{H}}||H|| + ||H||\mathrm{Tr}(A_{\mathcal{H}})^2 - 3||H||\mathrm{Tr}((\mathcal{A}^\phi_\xi)^2)_{\mathcal{H}}\right) g_{\mathcal{H}}(X,Y)
\end{aligned} \tag{5.3}$$

Also, we have

$$\begin{aligned}
(\nabla^{\mathcal{H}}_{\mathrm{grad}||H||}(S_{\varepsilon,\delta})_{\mathcal{H}})(X,Y) &= \frac{1}{||H||}(\nabla^{\mathcal{H}}_{\mathrm{grad}||H||} h_{\mathcal{H}})(X,Y)\\
&\quad -\frac{||\mathrm{grad}||H||\,||^2}{||H||^2} h(X,Y)\\
&\quad +\frac{3n^2L}{||H||^4}||\mathrm{grad}||H||\,||^2 g(X,Y)
\end{aligned} \tag{5.4}$$

and

$$\begin{aligned}
(\bar{\triangle}_{\mathcal{H}}(S_{\varepsilon,\delta})_{\mathcal{H}})(X,Y) &= \frac{1}{||H||}(\bar{\triangle}_{\mathcal{H}} h_{\mathcal{H}})(X,Y) - \frac{2}{||H||^2}(\nabla^{\mathcal{H}}_{\mathrm{grad}||H||} h_{\mathcal{H}})(X,Y)\\
&\quad +\frac{1}{||H||^3}\left(2||\mathrm{grad}||H||\,||^2 - ||H||\bar{\triangle}_{\mathcal{H}}||H||\right) h_{\mathcal{H}}(X,Y)\\
&\quad +\frac{3n^2L}{||H||^5}\left(-4||\mathrm{grad}||H||\,||^2 + ||H||\bar{\triangle}_{\mathcal{H}}||H||\right) g_{\mathcal{H}}(X,Y).
\end{aligned} \tag{5.5}$$

From (5.3), (5.4) and (5.5), we have

$$\begin{aligned}
\frac{\partial (S_{\varepsilon,\delta})_{\mathcal{H}}}{\partial t}(X,Y) &= \bar{\triangle}_{\mathcal{H}}(S_{\varepsilon,\delta})_{\mathcal{H}}(X,Y) + \frac{2}{||H||}(\nabla^{\mathcal{H}}_{\mathrm{grad}||H||}(S_{\varepsilon,\delta})_{\mathcal{H}})(X,Y)\\
&\quad +P(S_{\varepsilon,\delta})_{\mathcal{H}}(X,Y),
\end{aligned} \tag{5.6}$$

where $P(S_{\varepsilon,\delta})$ is defined by

$$\begin{aligned}P(S_{\varepsilon,\delta})(Z,W) := & -2((A_{\mathcal{H}})^2)_{\sharp}(Z,W) - 2((\mathcal{A}^{\phi}_{\xi})^2)_{\sharp}(Z,W) - \frac{1}{||H||}\mathcal{R}(Z,W)\\ & +\frac{1}{||H||^2}\left(2||H||\mathrm{Tr}((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}} + 4n^2L\right)h_{\mathcal{H}}(Z,W)\\ & +\frac{6n^2L}{||H||^3}\left(\mathrm{Tr}(A_{\mathcal{H}})^2 - 3\mathrm{Tr}((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}} + \frac{2||\mathrm{grad}\,||H||\,||^2}{||H||^2}\right)g_{\mathcal{H}}(Z,W)\\ & +\varepsilon g_{\mathcal{H}}(Z,W) - 2\varepsilon(\delta+t)||H||h_{\mathcal{H}}(Z,W)\end{aligned}$$

for $Z, W \in \pi_M^*TM$. Suppose that a positive number δ as in $(*)$ does not exist. Fix a sufficiently small positive number δ. For some $\varepsilon_0 > 0$, there exists $(x_0, t_0) \in M \times [0, \delta)$ such that $((S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)} > 0$ does not hold. Here we take t_0 as smally as possible. We have $\mathrm{Ker}((S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)} \neq \{0\}$ and $((S_{\varepsilon_0,\delta})^{\sharp}_t)_{\mathcal{H}_t} \geq 0$ $(\forall\, t \in [0, t_0])$. Take $X \in \mathrm{Ker}((S_{\varepsilon_0,\delta})_{\mathcal{H}})_{(x_0,t_0)})$ with $g(X, X) = 1$. Since

$$h(X,Y) = \left(\frac{2n^2L}{||H||^2} - \varepsilon_0(\delta+t_0)||H||\right)g(X,Y) \quad (\forall\, Y \in \mathcal{H}),$$

we have

$$A_{\mathcal{H}}X = \left(\frac{2n^2L}{||H||^2} - \varepsilon_0(\delta+t_0)||H||\right)X.$$

For simplicity, we set $\lambda_1 := \frac{2n^2L}{||H||^2} - \varepsilon_0(\delta + t_0)||H||$. By using the first relation in Lemma 3.10, we have

$$\begin{aligned}P(S_{\varepsilon_0,\delta})_{\mathcal{H}}(X,X) = & \frac{6n^2L}{||H||^3}\mathrm{Tr}(A_{\mathcal{H}})^2 + \frac{12n^2L}{||H||^5}||\mathrm{grad}||H||\,||^2\\ & -2((\mathcal{A}^{\phi}_{\xi})^2)_{\sharp}(X,X) - \left(\frac{14n^2L}{||H||^3} + 2\varepsilon_0(\delta+t_0)\right)\mathrm{Tr}((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}}\\ & +\frac{4}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{A}^{\phi}_X\bullet, \mathcal{A}^{\phi}_X(A_{\mathcal{H}}\bullet)\rangle - \frac{4}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{A}^{\phi}_{\bullet}X, \mathcal{A}^{\phi}_{\bullet}(A_{\mathcal{H}}X)\rangle\\ & +\frac{3}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_X\xi, \mathcal{A}^{\phi}_{\bullet}X\rangle - \frac{1}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle(\widetilde{\nabla}_X\mathcal{A}^{\phi})_{\bullet}\xi, \mathcal{A}^{\phi}_X\bullet\rangle\\ & +\frac{2}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle(\widetilde{\nabla}_{\bullet}\mathcal{A}^{\phi})_{\bullet}X, \mathcal{A}^{\phi}_X\xi\rangle + \varepsilon.\end{aligned} \tag{5.7}$$

Hence, since $\mathrm{Tr}(A_{\mathcal{H}})^2 \geq \frac{||H||^2}{n}$, $((\mathcal{A}^{\phi}_{\xi})^2)_{\sharp}(X,X) \leq 0$, $\mathrm{Tr}((\mathcal{A}^{\phi}_{\xi})^2)_{\mathcal{H}} \leq 0$ and the definition of L, we have

$$\begin{aligned}P(S_{\varepsilon_0,\delta})_{\mathcal{H}}(X,X) > & \frac{4}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{A}^{\phi}_X\bullet, \mathcal{A}^{\phi}_X(A_{\mathcal{H}}\bullet)\rangle\\ & -\frac{4}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle\mathcal{A}^{\phi}_{\bullet}X, \mathcal{A}^{\phi}_{\bullet}(A_{\mathcal{H}}X)\rangle.\end{aligned} \tag{5.8}$$

Since $A_{\mathcal{H}}X = \lambda_1 X$, $X \in \mathrm{Ker}(S_{\mathcal{H}})_{(u_0,t_0)}$ and $(S_{\mathcal{H}})_{(u_0,t_0)} \geq 0$, we see that λ_1 is the smallest eigenvalue of $A_{\mathcal{H}}$. Let $\{\lambda_i \,|\, i = 1, \cdots, n\}$ $(\lambda_1 \leq \cdots \leq \lambda_n)$ be the set of all eigenvalues of $A_{\mathcal{H}}$. Let $\{e_1, \cdots, e_n\}$ be an orthonormal base of $T_{(u_0,t_0)}M$ with $e_1 = X$ and $A_{\mathcal{H}}e_i = \lambda_i e_i$ $(i = 2, \cdots, n)$. Then we have

$$\begin{aligned}
&\frac{4}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle \mathcal{A}^{\phi}_{X}\bullet, \mathcal{A}^{\phi}_{X}(A_{\mathcal{H}}\bullet)\rangle - \frac{4}{||H||}\mathrm{Tr}^{\bullet}_{g_{\mathcal{H}}}\langle \mathcal{A}^{\phi}_{\bullet}X, \mathcal{A}^{\phi}_{\bullet}(A_{\mathcal{H}}X)\rangle \\
&= \frac{4}{||H||}\sum_{i=1}^{n}(\lambda_i - \lambda_1)\langle \mathcal{A}^{\phi}_{X}e_i, \mathcal{A}^{\phi}_{X}e_i\rangle \geq 0.
\end{aligned}$$

From (5.8) and this inequality, we obtain $P(S_{\varepsilon_0,\delta})_{\mathcal{H}}(X, X) \geq 0$. By using this fact, we can deduce a contradiction by imitating the discussion of (Step I) in the proof of Theorem 4.1. Hence we see that $(*)$ is true. Furthermore, by using $(*)$, we can show that $(S_{\mathcal{H}})_{(\cdot,t)} > 0$ holds for all $t \in [0, T)$ by imitating the discussion of (Step II) in the proof of Theorem 4.1. q.e.d.

6 Concluding remark

Let G/K be a compact Riemannian reductive homogeneous space whose isometry group G is compact and semi-simple, where we give G/K the metric induced from the Killing form B of the Lie algebra $\mathfrak{g}$ of G through the reductive decomposition of $\mathfrak{g}$. Here we note that G/K is of non-negative curvature. Let $H^0([0,1], \mathfrak{g})$ and $H^1([0,1], G)$ be as in Example of Section 3. This group $H^1([0,1], G)$ acts on $H^0([0,1], \mathfrak{g})$ isometrically and transitively as a gauge action. Set $P(G, \{e\} \times K) := \{g \in H^1([0,1], G) \,|\, (g(0), g(1)) \in \{e\} \times K\}$, where e is the identity element of G. This group $P(G, \{e\} \times K)$ acts on $H^0([0,1], \mathfrak{g})$ freely and isometrically, and the orbits of this action are minimal regularizable submanifolds. Also the orbit space of this action is isometric to G/K. Denote by ϕ the orbit map of this action, which is a Riemannian submersion. Let $\bar{M} (\subset G/K)$ be a hypersurface in G/K and $\bar{f}$ the inclusion map of $\bar{M}$ into G/K. Set $M := \phi^{-1}(\bar{M})$ and f the inclusion map of M into $H^0([0,1], \mathfrak{g})$. Then $f : M \hookrightarrow H^0([0,1], \mathfrak{g})$ is a $P(G, \{e\} \times K)$-invariant hypersurface. Let $\bar{f}_t$ $(0 \leq t < T)$ be the mean curvature flow starting from $\bar{f}$ and f_t $(0 \leq t < T)$ the regularizable mean curvature flow starting from f. Then we have $\bar{f}_t \circ \phi|_M = \phi \circ f_t$. Hence we can derive the strongly convexity preservability theorem by G. Huisken (see [Hu2, Theorem ?]) from Theorem 5.1. Here we note that Huisken [Hu2] treats the mean curvature flow in a complete Riemannian manifold in general.

References

[Ha] R. S. Hamilton, Three-manifolds with positive Ricci curvature, J. Differential Geom. **17** (1982) 255-306.

[Hu1] G. Huisken, Flow by mean curvature of convex surfaces into spheres, J. Differential Geom. **20** (1984) 237-266.

[Hu2] G. Huisken, Contracting convex hypersurfaces in Riemannian manifolds by their mean curvature, Invent. math. **84** (1986) 463-480.

[HLO] E. Heintze, X. Liu and C. Olmos, Isoparametric submanifolds and a Chevalley-type restriction theorem, Integrable systems, geometry, and topology, 151-190, AMS/IP Stud. Adv. Math. 36, Amer. Math. Soc., Providence, RI, 2006.

[Koi1] N. Koike, On proper Fredholm submanifolds in a Hilbert space arising from submanifolds in a symmetric space, Japan. J. Math. **28** (2002) 61-80.

[Koi2] N. Koike, Collapse of the mean curvature flow for equifocal submanifolds, Asian J. Math. **15** (2011) 101-128.

[O'N] B. O'Neill, The fundamental equations of a submersion, Michigan Math. J. **13** (1966) 459-469.

[P] R.S. Palais, Morse theory on Hilbert manifolds, Topology **2** (1963) 299-340.

[PT] R.S. Palais and C.L. Terng, Critical point theory and submanifold geometry, Lecture Notes in Math. **1353**, Springer, Berlin, 1988.

[T] C.L. Terng, Proper Fredholm submanifolds of Hilbert space, J. Differential Geometry **29** (1989) 9-47.

[TT] C.L. Terng and G. Thorbergsson, Submanifold geometry in symmetric spaces, J. Differential Geometry **42** (1995) 665-718.

[Z] X. P. Zhu, Lectures on mean curvature flows, Studies in Advanced Math., AMS/IP, 2002.

Department of Mathematics, Faculty of Science
Tokyo University of Science, 1-3 Kagurazaka
Shinjuku-ku, Tokyo 162-8601 Japan
(koike@ma.kagu.tus.ac.jp)